Volume Title
ASP Conference Series, Vol. **Volume Number**
Author
© **Copyright Year** Astronomical Society of the Pacific

arXiv:1402.1549v1 [astro-ph.GA] 7 Feb 2014

The Interstellar Reddening Law within 3kpc from the Sun

Hwankyung Sung,[1] and M. S. Bessell[2]

[1]*Department of Astronomy and Space Science, Sejong University, 209 Neungdongro, Kwangjin-gu, Seoul 143-747, Korea*

[2]*Research School of Astronomy and Astrophysics, Australian National University, MSO, Cotter Road, Weston, ACT 2611, Australia*

Abstract. We have investigated the interstellar reddening law of young open clusters within 3kpc from the Sun using optical, near-IR 2MASS, and *Spitzer* IRAC data. The total-to-selective extinction ratio R_V of 162 young open clusters (log $t_{age} \lesssim 7.3$) listed in the open cluster database WEBDA is determined from the color excess ratios. The young open clusters in the Sgr-Car arm show a relatively higher R_V, those in the Per arm and in the Cygnus region of the local arm show a relatively smaller value, and those in the Mon-CMa region of the local arm show a normal value ($R_V \approx 3.1$).

1. Introduction

To determine the physical parameters of stars and clusters accurately, we should know the correct interstellar reddening. Precise knowledge of R_V is of crucial importance for the determination of distances of reddened stars or clusters. Many investigators determine the total extinction A_V by assuming the total-to-selective extinction ratio R_V to be 3.1. In 1940s, from multicolor photometry of many O and B type stars Stebbins & Whitford (1943) concluded that the law of selective absorption is the same for all directions in the Galaxy. Later Johnson & Borgman (1963) conducted multi-color photometry of O, B stars and showed that there is no unique reddening law. In addition, they showed that there is a minor variation of R_V with galactic longitude. Later Serkowski et al. (1975) confirmed such a variation from the spatial variation of wavelength of maximum polarization. Whittet (1977) presented a functional form of R_V variation against galactic longitude. Recently Fitzpatrick & Massa (2009) showed that the interstellar reddening law R_V is different for different sightlines.

Recently Hur et al. (2012) showed that an abnormal reddenig law for the intracluster medium of the young open clusters in the η Carina nebula. However they obtained a fairly normal reddening law for the foreground stars. On the other hand, Sung et al. (2013a) found a nearly normal reddening law toward the young open cluster NGC 6231 from the optical to mid-IR photometric data. Now it is possible to study the reddening law toward young open clusters in a homogeneous way because the homogeneous near-IR 2MASS data as well as mid-IR *Spitzer* IRAC images are available for many young open clusters. In this paper we present a preliminary result from our systematic investigation of the reddening law.



Figure 1. Reddening law of Trumpler 16 from the optical to mid-IR. The reddening for a given color with respect to $E(B-V)$ is shown. The optical and near-IR data are from Hur et al. (2012) and 2MASS, respectively, while mid-IR photometry was obtained from the *Spitzer* IRAC images. The solid and dashed lines represent the normal reddening law for the foreground ($R_{V,fg} = 3.1$) and the abnormal reddening law for the intracluster medium of Trumpler 16 region ($R_{V,cl} = 4.6$), respectively. Large dots and squares denote stars with higher proper motion membership probability ($P_\mu \geq 70$) and with lower probability ($P_\mu < 30$), respectively.

2. Large Scale Variation of R_V

The reddening law for the young open clusters is determined as the same way as the method described in Hur et al. (2012); Sung et al. (2013b). The intrinsic color relations between $(B-V)_0$ and $(V-\lambda)_0$ for 2MASS JHK_s bands are presented in Sung et al. (2013b), and those for *Spitzer* IRAC bands will be published in the forthcoming paper.

We have analyzed the photometric data for 192 young open clusters with $\log \tau_{age} \lesssim 7.3$. Among them 16 clusters (e.g. Mayer 1, Be 65, Cz 13, Cr 173, etc) are either lack of UBV photoelectric or CCD data or $(U-B)$ color. And 4 clusters (e.g. Cz 20, NGC 2453, etc) are not young open clusters, 6 clusters (Cr 96, Cr 107, NGC 2448, etc) seem to be not clusters, and 4 clusters (BH 121, BH 205, IC 2948, and Loden 153) are parts of the other young clusters. And therefore we determined the reddening law of 162 young open clusters. We also determined the color excess ratios in the *Spitzer* IRAC bands for 38 clusters. The R_V of the young open clusters is determined the relation between R_V and color excess ratios by Guetter & Vrba (1989). Figure 1 shows the color excess ratios for the OB stars in the young open cluster Trumpler 16. A few foreground B type stars show a fairly normal reddening law of $R_{V,fg} = 3.1$, while cluster OB stars behave abnormally and well match to the solid line of $R_{V,cl} = 4.6$. Although most stars do not show any excess emission in the optical to near-IR, some stars show an evident excess emission in the mid-IR, especially in 8.0 μm.

In this paper we only describe the reddening law of the general interstellar medium. And therefore we only take the $R_{V,fg}$ for the young open clusters with an abnormal $R_{V,cl}$. Figure 2 shows the R_V with respect to galactic longitude. For some clusters we cannot



Figure 2. Plot of R_V against galactic longitude l. The solid curve is the variation of R_V from Whittet (1977). Large dots represent the R_V determined from the color excess ratios, while open circles denote the R_V determined from less reliable data. The size of symbols is proportional to the number of OB stars used in the determination of R_V.

determine the R_V reliably because of small values of reddening or no early type stars in the foreground. The young open clusters in the Sgr-Car arm show a relatively high value ($R_V \approx 3.2$), those in the Per arm and in the Cygnus region have a smaller value ($R_V \approx 2.9$), and those in the Mon-CMa regions have a nearly normal value ($R_V \approx 3.1$). In addition the fluctuation of R_V is also noticeable ($\sigma_{R_V} \approx 0.15$). However the overall variation of R_V well follows the equation given by Whittet (1977) except for the CMa-Vel direction.

3. Conclusion

From the analysis of photometric data for 162 young open clusters we confirmed the large scale variation of the reddening law in the galactic plane, which implies the real variation of dust size distribution in the galactic plane. We also found a large fluctuation of R_V for a given region. The cause of this fluctuation is still uncertain, and therefore a systematic photometric survey of open clusters to provide homogeneous photometric data, e.g. the Sejong Open cluster Survey (SOS) (Sung et al. 2013b) is required to confirm the real fluctuation.

Acknowledgments. H.Sung acknowledges the support of the National Research Foundation of Korea (NRF) funded by the Korea Government (MEST) (Grant No.2013031015).

References

Fitzpatrick, E. L., & Massa, D. 2009, ApJ, 699, 1209
Guetter, H. H., & Vrba, F. J. 1989, AJ, 98, 611
Johnson, H. L., & Borgman, J. 1963, BAN, 17, 115
Hur, H., Sung, H., & Bessell, M. S. 2012, AJ, 143, 41
Serkowski, K., Mathewson, D. S., & Ford, V. L. ApJ, 196, 261
Stebbins, J., & Whitford, A. E. 1943, ApJ, 98, 20
Sung, H., Sana, H., & Bessell, M. S. 2013, AJ, 145, 37
Sung, H., Lim, B., Bessell, M. S., KIm, J. S., Hur, H., Chun, M.-Y., & Park, B.-G. 2013, JKAS, 46, 103
Whittet, D. C. B. 1977, MNRAS, 180, 29